

RID® LICE KILLING SHAMPOO (With Conditioner)

Drug Facts

Active ingredients	Purpose
Piperonyl butoxide (4%).....	Lice treatment
Pyrethrum extract (equivalent to 0.33% pyrethrins).....	Lice treatment

Uses treats head, pubic (crab), and body lice

Warnings

For external use only

Do not use

- near the eyes
- inside the nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- **Important: Read warnings before use**

Adults and children 2 years and over:

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

Treat

- apply thoroughly to **DRY HAIR** or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for 10 minutes, but no longer
- use warm water to form a lather, shampoo, then thoroughly rinse
- for head lice, towel dry hair and comb out tangles

Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.



RID® LICE KILLING SHAMPOO (With Conditioner)

- pin back each strand of hair after combing
 - clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
 - after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
 - check daily for any lice/nits that you missed
 - a second treatment must be done in 7 to 10 days to kill any newly hatched lice
 - if infestation continues, see a doctor for other treatments
- Children under 2 years:** ask a doctor

Other information

- keep carton for important product information
- see Consumer Information Insert for additional information
- avoid excessive heat

Inactive ingredients ammonium laureth sulfate, dehydrated alcohol, fragrance, isopropyl alcohol, PEG-25 hydrogenated castor oil, polyquaternium-10, purified water

Questions or comments? 1-800-RID-LICE (1-800-743-5423)
(Mon – Fri 9AM – 8PM EST) www.ridlice.com