



# 3-STEP PROCESS FOR LICE TREATMENT<sup>1</sup>

See carton for complete product information.

## Step 1 • Treat

### Apply RID<sup>®</sup> Lice Killing Shampoo

**Note: Do not treat** unless live lice or eggs have been found on the individual's head.



**ACTIVE INGREDIENTS:**

Piperonyl butoxide 4%  
Pyrethrum extract  
(equivalent to 0.33% pyrethrins)

**INACTIVE INGREDIENTS:**

ammonium laureth sulfate, dehydrated alcohol, fragrance, isopropyl alcohol, PEG-25 hydrogenated castor oil, polyquaternium-10, purified water

**TREATS:**

Head, pubic (crab), and body lice.

**WARNINGS:**

**For External use only**

- Do not use near the eyes, inside nose, mouth, vagina, or on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.
- Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthmatic attack.
- When using this product keep eyes tightly closed and protect eyes with a washcloth or towel. If product gets in eyes, flush with water right away. Scalp itching or redness may occur.
- Stop use and consult a doctor if breathing difficulty occurs, eye irritation occurs, skin or scalp irritation continues or infection occurs.
- **Keep out of reach of children!** swallowed, get medical help or contact a Poison Control Center right away.



**A** Use towels to protect eyes and clothes from treatment. Apply **RID<sup>®</sup> Lice Killing Shampoo** to **DRY HAIR** or affected area. Apply enough product to saturate. Thoroughly massage product into scalp, behind ears and onto back of neck.



**B** Allow product to remain on hair for 10 minutes, but no longer.



**C** Add warm water and massage to form lather. Rinse thoroughly, e.g., in a sink.

**D** Repeat this step in 7-10 days to kill any newly hatched lice.

## Step 2 • Remove

### Comb out lice and eggs with RID<sup>®</sup> Lice Comb or other fine-tooth comb and RID<sup>®</sup> Comb-Out Spray

**Note:** RID<sup>®</sup> Lice Comb and Comb-Out Spray are not included in RID<sup>®</sup> Shampoo-only products.



**Use:** RID<sup>®</sup> Lice & Egg Comb-Out Spray detangles hair in order to make lice and egg comb-out easier.

**Ingredients:** Water, Glycerin, Propylene Glycol, Polysorbate 20, Cetrimonium Chloride, Amodimethicone, Phenoxyethanol, Fragrance, DL Panthenol, Trideceth-12, Methylchloroisothiazolinone, Methylisothiazolinone.

**Warnings:** For external use only. Keep out of reach of children. Avoid contact with eyes.



**A** Apply a small amount of **RID<sup>®</sup> Comb-Out Spray** and use a wide-tooth comb to remove any tangles.



**B** Lift a 1-2" section of hair. Starting as close to scalp as possible, slowly comb section all the way to ends of hair with **RID<sup>®</sup> Lice Comb**, using additional Comb-Out Spray to remove tangles as needed. Use a tissue or paper towel to wipe any lice or eggs off comb.



**C** Pin back this section of hair. Continue to comb out one 1-2" section at a time until entire head has been combed and pinned.



**D** Remove pins and rinse thoroughly with warm water. You may use your regular shampoo now if desired. The RID<sup>®</sup> Lice Comb should be disinfected for reuse by soaking in hot water (above 130°F) for 10 minutes.

## Step 3 • Control

### Treat the Home

**Note:** RID<sup>®</sup> Home Spray is not included in RID<sup>®</sup> Shampoo-only products or in the RID<sup>®</sup> Lice Elimination Essentials Kit.



Use **RID<sup>®</sup> Home Lice, Bedbug & Dust Mite Spray** to kill lice and eggs on unwashable items such as rugs, furniture or car interiors. Do not use RID<sup>®</sup> Home Lice, Bedbug and Dust Mite Spray on the head or body.

## Additional Tips for Success

- Be sure to complete all three steps in the RID<sup>®</sup> Lice Treatment process. This includes a second shampooing between 7-10 days after the initial treatment, but no sooner. If any eggs were not initially combed out, lice may hatch and can be killed with the second treatment. Newly hatched lice are not mature enough to lay new eggs.
- Carefully inspect infested family members daily for at least 2 weeks after the second shampooing. If you see eggs or nits, consult your doctor.
- Teach children not to share personal items, such as hats, combs and brushes and to avoid prolonged contact with a person who has lice.

**Questions?**  
Call the RID<sup>®</sup> Expert Support help line at 1-800-RID-LICE (1-800-743-5423).